



Dermatology Reports

<https://www.pagepress.org/journals/index.php/dr/index>

eISSN 2036-7406



SIDCO

Società Italiana di Dermatologia
Chirurgica, Oncologica, Correttiva ed Estetica

Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. **Dermatology Reports** is, therefore, E-publishing PDF files of an early version of manuscripts that undergone a regular peer review and have been accepted for publication, but have not been through the copyediting, typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear on a regular issue of the journal.
E-publishing of this PDF file has been approved by the authors.

Please cite this article as: Tchernev G, Kordeva S. Modified advancement transposition flap for squamous cell carcinoma with periauricular location. Dermatol Rep 2024 [Epub Ahead of Print] doi: 10.4081/dr.2024.9977



© the Author(s), 2024
Licensee PAGEPress, Italy

Submitted: 26/02/2024 – Accepted 15/03/2024

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Modified advancement transposition flap for squamous cell carcinoma with periauricular location

Georgi Tchernev,^{1,2} Simona Kordeva¹

¹Onkoderma – Clinic for Dermatology, Venereology and Dermatologic Surgery, Sofia; ²Department of Dermatology and Venereology, Medical Institute of Ministry of Interior, Sofia, Bulgaria

Correspondence: Simona Kordeva, Onkoderma- Clinic for Dermatology, Venereology and Dermatologic Surgery, General Skobelev 26, 1606 Sofia, Bulgaria.

Tel.: +359.884959176.

E-mail: simonakordeva97@gmail.com

Conflict of interest: the authors declare no potential conflict of interest.

Consent for publication: the patient gave his consent to use his personal data for the publication of this case report and any accompanying images.

The case

A 74-year-old male presented to the dermatology department with tumor formation located in the left auricle area, measuring 2 cm in size. The tumor exhibited surface erosions and periphereal erythema. Additionally, in the right lower eyelid and temporal regions, clinically suspected lesions for squamous cell carcinomas were noted. The patient had a previous history of 15 surgical interventions in the facial area, primarily for squamous cell carcinomas, with one documented removal of a basal cell carcinoma. A clinical diagnosis of squamous cell carcinoma was established, and the patient was recommended surgical excision under local anesthesia.

How would you remove the lesion?

Figure 1

Our choice

The tumor formation in the periauricular region was removed with an elliptical excision with a surgical safety margin of 4 mm in all directions. Primary closure with single interrupted sutures or secondary wound healing were not an option due to the size and location of the primary defect. The generated tension will compromise the anatomical integrity of the area, resulting in both functional impairments and aesthetic issues. We also took into consideration the close proximity to the superficial temporal vessels.

Considering the size of the primary defect and the patient's skin elasticity, our team decided to utilize a modified advancement transposition flap to repair the large defect while also preserving the anatomical integrity of the area. The flap will efficiently redistribute the primary tension vectors generated during defect closure, resulting in a more refined outcome. Additionally, the flap tissue will have similar vascular supply and innervation.

The flap was formed in the upper part of the primary defect, following careful dissection and undermining down to the hypodermis. The pedicle will consist of the subcutaneous tissue beneath the flap. Subsequently, the flap was transposed towards the jaw direction. The superficial temporal vessels were successfully preserved. The skin edges were adapted using 1-00 sutures, extending from the orbit to the ear and from the nose to the periauricular area, minimizing the tension vectors and evenly distributing it in all directions. (Figure 2a-c)

The sutures were removed 14 days post-surgery. The flap remained healthy without any postoperative complications. After the 1-month follow-up, the patient exhibit improvement, and an aesthetic outcome was achieved.

Comment

Primary wound defects that extend beyond the limits of primary closure should be considered for reconstruction using a local advancement flap (1). When employing a flap reconstructive technique, several important factors should be considered, including the size and location of the primary defect, the primary tension vectors and their distribution, potential impact on adjacent anatomical structures, flap vascularity and perfusion, and etc (1). Local flaps are “ideal” candidates when the adjacent tissue is healthy or unaffected by the malignant process, allowing for immediate tissue relocation to address a nearby primary defect (1).

With the use of a modified advancement flap we ensured minimal trauma to the surrounding tissues while also achieving a more aesthetically refined final look. The vascular supply was effectively preserved, and the postoperative period proceeded without any complications.

The outcome

Figure 3 – a) 14 days post-surgery b) 1 month follow-up

References

1. Etzkorn JR, Zito PM, Hohman MH, et al. Advancement Flaps. [Updated 2024 Jan 29]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK431081/>



Figure 1.



Figure 2.



Figure 3.