How does the *NPM1* mutant induce leukemia?

Paolo Sportoletti

Hematology and Clinical Immunology
Section, Department of Clinical
and Experimental Medicine,
University of Perugia, Perugia, Italy

Abstract

NPM1 is the most frequently mutated gene in AML and the role of the *NPM1* mutant in acute myeloid leukemia along with its leukemogenic potential are still under investigation.

NPM1 genetic alterations can contribute to leukemogenesis through the direct oncogenic effect of the mutant protein and the concomitant loss of one functional allele. *Npm1* loss determines tumor development in the mouse while in human *NPM1* maps in a chromosomal region frequently lost in myelodysplastic syndrome (MDS). The *NPM1* mutant cytoplasmic delocalization in leukemic blasts alters multiple cellular pathways through either loss or gain of function effects on different protein partners.

Here we discuss the most relevant studies on the role of the *NPM1* molecule in hematological malignancies and both *in vitro* and *in vivo* studies that are trying to elucidate the way by which the *NPM1* mutation induces leukemia.

Introduction

NPM1 is a multifunctional phosphoprotein that encodes for a number of functional domains through which the molecule is able to bind many partners in distinct cellular compartments.¹ *NPM* displays nucleolar localization and constantly shuttles between the nucleus and the cytoplasm. The shuttling activity of *NPM1* along with its capacity to interact with many partners involves the protein in multiple cellular functions.

These functions include ribosome biogenesis and the transport of pre-ribosomal particles, maintenance of genomic stability through the control of cellular ploidy and centrosome duplication and the control of cellular proliferation. *NPM1* is also a survival factor and participates in DNA-repair processes. In addition, *NPM1* is involved in regulating the activity and stability of crucial tumor suppressors such as ARF and p53. The *NPM1* polypeptide chain has a modular structure containing distinct sequence motifs including a nuclear-export signal (NES) and a nucleolar localization

signal (NuLS) that resides in the C-terminal region of the protein. In acute myeloid leukemia (AML) patients the NuLS is the target of the C-terminal mutation that determines its substitution into an extra NES, thus resulting in an aberrantly localized protein that resides in the cytoplasm of leukemic blasts.² The shuttling activity of *NPM1* and its proper sub cellular localization are thought to be crucial for normal cellular homeostasis and the identification of the *NPM1* mutant counterpart in AML patients has emphasized this hypothesis. Thus, genetic alteration of *NPM1* can contribute to oncogenesis by directly affecting *NPM1* functions.

NPM1 has been directly implicated in human cancer.¹ In fact the *NPM1* protein is overexpressed in solid tumours of different origin like gastric, colon, ovarian and prostate carcinomas. *NPM1* is one of the most frequent targets of genetic alterations in haematopoietic tumours. It is found translocated with distinct partner genes in several diseases such as acute promyelocytic leukemia (APL), anaplastic large cell lymphoma (ALCL), acute myeloid leukemia (AML) and myelodysplasia. Moreover, the region of chromosome 5 that *NPM1* maps to, is a genetic target of deletions in both *de novo* and therapy related MDS, a pre-leukemic condition with ineffective production of myeloid blood cells. More recently *NPM1* has been found mutated and aberrantly localized in the cytoplasm of myeloid blasts in a high proportion of AML patients with normal karyotype.² Although the *NPM1* gene was strongly implicated in cancer pathogenesis, how the cytoplasmic localized *NPM1* mutant protein (*NPMc+*) promotes leukemia is still under investigation.

NPM1 gene alterations contributing to leukemogenesis

The alteration of *NPM1* in human cancer can lead to tumor development as a result of two major mechanisms involving both the presence of a mutated product and the reduction of the wild type *NPM1* dosage and level of expression. Indeed, *NPM1* gene alterations lead to the generation of mutated products associated with hematopoietic malignancies (in the case of AML the *NPMc+* mutant), and to the concomitant loss of one functional allele of the gene leading to *NPM1* heterozygosity.

The reduction of the *NPM* dosage to heterozygosity has been shown to lead to altered *NPM1* functions and tumour susceptibility³ indicating that normal levels of expression are required for the cell to prevent tumor development. Moreover mutated products retain the ability to heterodimerize with the residual wild-type protein, and they potentially interfere with its functions. In particular the dominant-negative effect of the mutated *NPM* determines the delocalization of the remaining

Correspondence: Paolo Sportoletti, Hematology and Clinical Immunology Section, Department of Clinical and Experimental Medicine, University of Perugia, Perugia, Italy.
E-mail: sportolp@gmail.com

Key words: *NPM1*, leukemia.

Received for publication: 4 May 2011.
Accepted for publication: 4 June 2011.

This work is licensed under a Creative Commons Attribution NonCommercial 3.0 License (CC BY-NC 3.0).

©Copyright P. Sportoletti, 2011
Licensee PAGEPress, Italy
Pediatric Reports 2011; 3(s2):e6
doi:10.4081/pr.2011.s2.e6

wild-type *NPM* protein and exacerbates the *NPM* functional loss. On the other end there is the capability for the *NPMc+* mutant itself to function as activated oncogene. Since the *NPM1* mutation always results in aberrant cytoplasmic dislocation of the mutant protein, this event appears critical for leukemogenesis. The increased *NPM1* export into the cytoplasm can affect multiple cellular pathways and drive leukemia by either a loss-of-function^{4,5} or a gain-of-function.⁶ In fact, the *NPMc+* nucleolar interactors can be delocalized by the mutant into leukemic cells cytoplasm and their activity can be significantly impaired. On the other end the *NPMc+* mutant may acquire the capability to interact with new protein partners in the cytoplasm. All these activities can lead to leukemia and there are a number of *in vitro* and *in vivo* data supporting this hypothesis that will be analyzed in the next paragraphs.

The role of *Npm* functional loss in tumor development

Data on the role of *Npm* functional loss in tumor development arise from mouse work that has been done in the Pandolfi's lab in the last five years. *Npm* complete knockout resulted in embryonic lethality due to developmental abnormalities of the brain and of the hematopoietic system.³ In this setting *Npm* inactivation was able to determine cancer susceptibility *in vitro*. The other main observation of this study implicated *Npm* in the pathogenesis of human MDS. *Npm* heterozygous mice showed some of the features found in the human syndrome. Peripheral blood showed significantly increased MCV and RDW values while the bone marrow presented a high proportion of dysplastic erythroid precursors and megakaryocyte dyspoiesis. This MDS status evolved into overt leukemia as in humans, demonstrating that *Npm* act as tumor suppressor gene *in vivo*.⁷ In fact, *Npm1* heterozy-

gous mice displayed higher susceptibility to develop tumor than their wild type counterparts (29% vs 6,3%). Pathological analysis revealed that 75% of the diseased mice were affected by hematological malignancies, including myeloid leukemia, while no abnormalities were detected in the hematopoietic organs of wild type animals analyzed. Npm loss of function at the centrosome has been involved in tumor development in these mice as bone marrow cytopins from blasts of affected mice clearly showed an increased number of cells displaying multiple centrosomes.

Is NPMc+ an oncogene *in vitro*?

Until very recently there was no published evidence that NPMc+ would transform primary cells *in vitro*. At first, the NPMc+ oncogenic capacity *in vitro* has been debated as a study suggested a tumor suppressive role for the NPMc+ mutant because its overexpression in ARF null cells end up inhibiting cell proliferation.⁸

In order to establish whether NPMc+ is an oncogene *in vitro*, the effect of NPMc+ overexpression in primary MEFs has been studied performing classical oncogenic transformation assays in soft agar.⁹ The study demonstrated that NPMc+ is able to transform when it is combined with E1A. E1A is an adenoviral oncogene that induces a E2F1-dependent cell cycle progression. At the same time it induces apoptosis through Arf and p53. NPMc+ blocks Arf, allowing E1A to evade the p53 induced apoptosis in transformed cells while the addition of E1A completely abolishes the NPMc+ induced senescent phenotype and allows transformation. This represented the first demonstration that NPMc+ is an oncogene *in vitro* and suggested the need for a cooperative event in leukemogenesis.

Data arising from other studies described a model in which mutations of NPM1 seem to simultaneously dampen a tumor-suppressor pathway⁴ (p53–ARF) and enhance an oncogenic one (MYC).⁵ In fact, NPMc+ was demonstrated to be able to reduce the ability of ARF to initiate a p53 response and to induce cell cycle arrest. Bonetti *et al.* show that, in the absence of NPM or in the presence of NPMc+, FBW7 γ loses its nucleolar localization and is rapidly degraded by the proteasome. FBW7 γ is a nucleolar ubiquitin ligase previously implicated in the ubiquitination/degradation of MYC. NPMc+ maintains the property of interacting with FBW7 γ but delocalizes it to the cytoplasm, where it is degraded, thus leading to accumulation of MYC and increased MYC signaling.

The NPM1 mutant may also exert its transforming properties through gain-of function in cytoplasm.⁶ Interestingly, the NPM1 mutant binds caspase 6 and 8 in the cytoplasm and specifically inhibits the activities of these cell-death caspases through direct interaction with their cleaved active forms. Moreover NPMc+ is

able to suppress caspase 6 and 8-mediated myeloid differentiation.

Is NPMc+ an oncogene *in vivo*?

Until very recently there was no published data on NPMc+ oncogenic activity *in vivo*. However, cytoplasmic mutated NPM has been retained for eight years in a xenotransplant model of NPMc+ acute myeloid leukemia in immunodeficient mice by subcutaneous injection of AML blasts from a patient.¹⁰ This study demonstrated that cytoplasmic mutated NPM1 is stable in AML and suggested that the NPM1 mutation represent a founder genetic lesion in AML.

In an attempt to understand the *in vivo* role of NPMc+ in AML, it has been recently developed a model showing an impact of the mutation on myelopoiesis with no progression toward overt leukemia.¹¹ NPMc+ was overexpressed under the control of the MRP8 promoter which lead the expression of NPMc+ in the common myeloid progenitors as well as mature granulocytes and monocytes. These mice came down with myeloproliferative disorders with splenomegaly and increased number of Gr1Mac-1 mature myeloid cells in the bone marrow and spleen. However, the latency was long and the phenotype was not fully penetrant among the different line analyzed and, more importantly, the NPM1 mutant alone was not able to initiate AML. Consequently, to exert its oncogenic effect, NPM1 may need to act under different conditions, such as targeting a specific stem cell/myeloid precursor and/or achieving a mutant to wild-type expression ratio that is appropriate for cytoplasmic delocalization of both nucleophosmin forms and/or being accompanied by a secondary cooperating event. However, the fact that NPMc+ is a bona fide oncogene *in vivo* comes from another interesting finding that NPMc+ also drives epithelial cancer given the fact that MRP8 is a leaky promoter in skin. Considering MRP8 is leaky in the keratinocytes, this presumably demonstrates that NPMc+ is an oncogene *in vivo*.

Data on zebrafish completely confirm what shown in the MRP8-NPMc+ transgenic mouse model.¹² In zebrafish, ubiquitous mutant NPMc+ not only caused expansion of primitive myeloid cells, but also resulted in increased numbers of definitive erythro-myeloid progenitors (*gata1*+/*lmo2*bright) and hematopoietic stem cells (*c-myb*+/*cd41*+) in the aorta ventral wall. In the zebrafish embryo, follow-up for AML development was not possible due to the transient nature of mutant NPM1 expression.

More recently a conditional knock-in allele of the NPM1 mutation A demonstrated that NPMc+ causes acute myeloid leukemia in mice.¹³ NPMc+ heterozygous activation in the hematopoietic system was associated with expansion of mature bone marrow mature myeloid cells as previously described. Knock-in

mutant display increased incidence of acute myeloid and lymphoid leukemias arising with a long latency suggesting the requirement for additional mutations. Insertional mutagenesis screens *in vivo* accelerated leukemia onset confirming the need for cooperative events in NPMc+ leukemogenesis.

Conclusions

The potential role of the NPM mutant as oncogene as well as the aberrant mechanism that leads to leukaemogenesis in NPMc+ AML is still under investigation. Here we have presented the most relevant studies describing how NPM1 alterations could contribute to leukemia. NPM1 expression and gene integrity are frequently altered in human hematological malignancies. Thus, the pathogenesis of NPMc+ AML has been proposed to depend on both NPM functional loss and the NPMc+ mutant oncogenic activity.

Genetic and molecular evidence has shed light on the mechanisms of NPM-mediated tumor suppression, demonstrating that Npm is a haploinsufficient suppressor of both myeloid and lymphoid malignancies in the mouse. NPMc+ is able to exert oncogenic potential by either loss or gain of function effects that impact on multiple cellular pathways. It has been demonstrated that NPMc+ act as a bona fide proto-oncogene *in vitro* and *in vivo* and suggested that additional cooperative genetic hits are required for leukemogenesis. Knock-in mouse models mimicking human NPM1-mutated AML are still under development and will be essential tools in the discovery of novel NPMc+ leukemogenic activities.

References

1. Grisendi S, Mecucci C, Falini B, Pandolfi PP. Nucleophosmin and cancer. *Nat Rev Cancer* 2006;6:493-505.
2. Falini B, Mecucci C, Tiacci E, et al. Cytoplasmic nucleophosmin in acute myelogenous leukemia with a normal karyotype. *N Engl J Med* 2005;352:254-66.
3. Grisendi S, Bernardi R, Rossi M, et al. Role of nucleophosmin in embryonic development and tumorigenesis. *Nature* 2005;437:147-53.
4. Colombo E, Martinelli P, Zamponi R, et al. Delocalization and destabilization of the Arf tumor suppressor by the leukemia-associated NPM mutant. *Cancer Res* 2006; 66:3044-50.
5. Bonetti P, Davoli T, Sironi C, et al. Nucleophosmin and its AML-associated mutant regulate c-Myc turnover through Fbw7 gamma. *J Cell Biol* 2008;182:19-26.

6. Leong SM, Tan BX, Bte Ahmad B, et al. Mutant nucleophosmin deregulates cell death and myeloid differentiation through excessive caspase-6 and -8 inhibition. *Blood* 2010;116:3286-96.
7. Sportoletti P, Grisendi S, Majid SM, et al. Npm1 is a haploinsufficient suppressor of myeloid and lymphoid malignancies in the mouse. *Blood* 2008;111:3859-62.
8. den Besten W, Kuo ML, Williams RT, Sherr CJ. Myeloid leukemia-associated nucleophosmin mutants perturb p53-dependent and independent activities of the Arf tumor suppressor protein. *Cell Cycle* 2005;4:1593-8.
9. Cheng K, Grisendi S, Clohessy JG, et al. The leukemia-associated cytoplasmic nucleophosmin mutant is an oncogene with paradoxical functions: Arf inactivation and induction of cellular senescence. *Oncogene* 2007;26:7391-400.
10. Falini B, Martelli MP, Mecucci C, et al. Cytoplasmic mutated nucleophosmin is stable in primary leukemic cells and in a xenotransplant model of NPMc+ acute myeloid leukemia in SCID mice. *Haematologica* 2008;93:775-9.
11. Cheng K, Sportoletti P, Ito K, et al. The cytoplasmic NPM mutant induces myeloproliferation in a transgenic mouse model. *Blood* 2010;115:3341-5.
12. Bolli N, Payne EM, Grabher C, et al. Expression of the cytoplasmic NPM1 mutant (NPMc+) causes the expansion of hematopoietic cells in zebrafish. *Blood* 2010;115:3329-40.
13. Vassiliou G, Cooper JL, Rad R, et al. A conditional knock-in allele of the type a cytoplasmic nucleophosmin mutation cooperates with a novel sleeping beauty transposon to cause acute myeloid leukaemia in mice. *Haematologica* 2010; 95(s2):240, abstr. 0578.

non commercial use only